## Human CD4/LEU3 Protein

#### Cat. No. CD4-HM204



Description	
Source	Recombinant Human CD4/LEU3 Protein is expressed from HEK293 with hFc at the C-Terminus.
	It contains Lys26-Trp390.
Accession	P01730
Molecular Weight	The protein has a predicted MW of 67.5 kDa. Due to glycosylation, the protein migrates to 70-80 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and	Storage
Committee	Lyophilized from 0.22µm filtered solution in PBS, 100mM L-arginine (pH 7.4). Normally 8% trehalose is added as


Formulation	Lyophilized from 0.22µm filtered solution in PBS, 100mM L-arginine (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

# **Background**

CD4, also known as L3T4, T4, and W3/25, is an approximately 55 kDa type I transmembrane glycoprotein that is expressed predominantly on thymocytes and a subset of mature T lymphocytes. It is a standard phenotype marker for the identification of T cell populations. Integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class II molecule:peptide complex.

## **Assay Data**

#### **Bis-Tris PAGE**




Human CD4 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

**SEC-HPLC** 




## **Assay Data**



The purity of Human CD4 is greater than 95% as determined by SEC-HPLC.

#### **ELISA Data**

Human CD4, hFc Tag ELISA 0.1µg Human CD4, hFc Tag Per Well



Log Biotinylated Anti-CD4 Antibody, hFc-Avi Tag Conc.(µg/ml)

Immobilized Human CD4 at 1µg/ml (100µl/Well) on the plate. Dose response curve for Biotinylated Anti-CD4 Antibody, hFc Tag with the EC50 of 15.4ng/ml determined by ELISA (QC Test).