Mouse IL-2 R alpha/CD25 Protein

Cat. No. CD5-MM125



Description	
Source	Recombinant Mouse IL-2 R alpha/CD25 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Glu22-Lys236.
Accession	P01590
Molecular Weight	The protein has a predicted MW of 25.76 kDa. Due to glycosylation, the protein migrates to 50-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

IL-2 receptor alpha (IL-2 R alpha), also known as Tac antigen and as CD25, was initially identified as a 55 kDa membrane glycoprotein that is capable of binding IL2.eceptor for interleukin-2. The receptor is involved in the regulation of immune tolerance by controlling regulatory T cells (TREGs) activity. TREGs suppress the activation and expansion of autoreactive T-cells.


Assay Data

Bis-Tris PAGE



Mouse IL-2 R alpha on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Mouse IL-2 R alpha is greater than 95% as determined by SEC-HPLC.


KAGTUS

Assay Data

ELISA Data


Mouse IL-2 R alpha, His Tag ELISA

0.5μg Mouse IL-2 R alpha, His Tag Per Well



Log Biotinylated Mouse IL-2, His-Avi Tag Conc.(μg/ml)

SPR Data



Immobilized Mouse IL-2 R alpha, His Tag at $5\mu g/ml$ (100 $\mu l/well$) on the plate. Dose response curve for Biotinylated Mouse IL-2, His Tag with the EC50 of 0.15 $\mu g/ml$ determined by ELISA (QC Test).

Mouse IL-2 R alpha, His Tag immobilized on CM5 Chip can bind Mouse IL-2, His Tag with an affinity constant of 30.49 nM as determined in SPR assay (Biacore T200).