Biotinylated Rhesus macaque CD5 Protein

lymphoma.


Cat. No. CD5-RM405B



Description	
Source	Recombinant Biotinylated Rhesus macaque CD5 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Arg25-Pro375.
Accession	F6V5Q9
Molecular Weight	The protein has a predicted MW of 41.7 kDa. Due to glycosylation, the protein migrates to 55-60 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per ug by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	CD5: a type I transmembrane protein found on T cells, thymocytes, and some B cells that is a ligand for CD72 and is involved in cellular activation or adhesion; expressed in B-cell chronic lymphocytic leukemia and T-cell

Assay Data

Bis-Tris PAGE



Biotinylated Rhesus macaque CD5 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


ELISA Data

Assay Data



Biotinylated Rhesus macaque CD5, His Tag ELISA

0.1μg Biotinylated Rhesus macaque CD5, His Tag Per Well



Log Anti-CD5 Antibody, hFc Tag Conc.(µg/ml)

Immobilized Biotinylated Rhesus macaque CD5, His Tag at $1\mu g/ml$ ($100\mu l/Well$) on streptavidin ($5\mu g/ml$) precoated plate. Dose response curve for Anti-CD5 Antibody, hFc Tag with the EC50 of 9.3ng/ml determined by ELISA.