

Cynomolgus CD27 Ligand/CD70 Protein

Cat. No. CD7-CM170




Description	
Source	Recombinant Cynomolgus CD27 Ligand/CD70 Protein is expressed from HEK293 with His tag at the N-Terminus. It contains Gln39-Pro194.
Accession	G7PYU6-1
Molecular Weight	The protein has a predicted MW of 18.4 kDa. Due to glycosylation, the protein migrates to 55-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per ug by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
CD70, also named CD27 ligand (CD27L), is a type II transmembrane glycoprotein belonging to the TNF superfamily (TNFSF) and has been designated TNFSF7. CD70 is a cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells.	

Assay Data

Bis-Tris PAGE




Cynomolgus CD27 Ligand on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

ELISA Data

Assay Data

Cynomolgus CD27 Ligand, His Tag ELISA
0.2µg Cynomolgus CD27 Ligand, His Tag Per Well



Immobilized Cynomolgus CD27 Ligand, His Tag at 2µg/ml (100µl/well) on the plate. Dose response curve for Cynomolgus/Rhesus macaque CD27, hFc Tag with the EC50 of 100.2ng/ml determined by ELISA (QC Test).