

Human CD79B Protein

Cat. No. CD7-HM19B



Description

Source	Recombinant Human CD79B Protein is expressed from HEK293 with His tag at the C-terminus.
	It contains Ala29-Asp159.
Accession	P40259-1
Molecular Weight	The protein has a predicted MW of 16.82 kDa. Due to glycosylation, the protein migrates to 35-50 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Supplied as 0.22 µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

CD79B (also known as B29, Ig beta and B cell antigen receptor complex-associated protein beta-chain) is a 36-40 kDa member of the Ig-Superfamily. It is required in cooperation with CD79A for initiation of the signal transduction cascade activated by the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. Enhances phosphorylation of CD79A, possibly by recruiting kinases which phosphorylate CD79A or by recruiting proteins which bind to CD79A and protect it from dephosphorylation.


Assay Data

Bis-Tris PAGE



Human CD79B on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Human CD79B is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Human CD79B, His Tag ELISA

0.1µg Human CD79B, His Tag Per Well



Immobilized Human CD79B, His Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Anti-CD79B Antibody, hFc Tag with the EC50 of 1.8ng/ml determined by ELISA.