

Rhesus macaque Complement Factor D/CFD Protein



Cat. No. CFD-CM101


Description	
Source	Recombinant Rhesus macaque Complement Factor D/CFD Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Gln21-Ala253.
Accession	H9EXC1-1
Molecular Weight	The protein has a predicted MW of 25.7 kDa. Due to glycosylation, the protein migrates to 26-30 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in 50mM Tris, 150mM NaCl (pH 8.0). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
Complement factor D is a serine protease essential for the activation of the alternative pathway and is expressed in the kidney, adipocytes, and macrophages. Factor D is found at relatively high levels in glomeruli suggesting that this component of the complement cascade could influence renal pathophysiology. Complement factor D or alternative pathway activation is needed to prevent spontaneous accumulation of C3 and IgM deposits within the mesangium.	

Assay Data

Bis-Tris PAGE



Rhesus macaque Complement Factor D on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Rhesus macaque Complement Factor D/CFD Protein

Cat. No. CFD-CM101



Assay Data




The purity of Rhesus macaque Complement Factor D is greater than 95% as determined by SEC-HPLC.

ELISA Data

Rhesus macaque Complement Factor D, His Tag ELISA

0.1µg Rhesus macaque Complement Factor D, His Tag Per Well



Immobilized Rhesus macaque Complement Factor D, His Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Anti-Complement Factor D Antibody, hFc Tag with the EC50 of 16.3ng/ml determined by ELISA.