

Biotinylated Human DKK1 N terminal Domain Protein



Cat. No. DKK-HM50NB


| Description      |   |
|------------------|---|
| Source           | Recombinant Biotinylated Human DKK1 N terminal Domain Protein is expressed from HEK293 with hFc tag and Avi tag at the C-Terminus.<br>It contains Thr32-Asp142. |
| Accession        | O94907  |
| Molecular Weight | The protein has a predicted MW of 40.25 kDa. Due to glycosylation, the protein migrates to 50-60 kDa based on Bis-Tris PAGE result.                             |
| Endotoxin        | Less than 1 EU per µg by the LAL method.  |
| Purity           | > 95% as determined by Bis-Tris PAGE<br>> 95% as determined by HPLC   |

| Formulation and Storage |   |
|-------------------------|---|
| Formulation             | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.  |
| Reconstitution          | Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.   |
| Storage                 | -20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

| Background  |  |
|---|--|
| Dickkopf-1 (Dkk1), the founding and best-studied member of the Dkk family, functions as an antagonist of canonical Wnt/β-catenin. Dkk1 is considered to play a broad role in a variety of biological processes. |  |

Assay Data

Bis-Tris PAGE



Biotinylated Human DKK1 N terminal Domain on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Biotinylated Human DKK1 N terminal Domain Protein

Cat. No. DKK-HM50NB



Assay Data



The purity of Biotinylated Human DKK1 N terminal Domain is greater than 95% as determined by SEC-HPLC.