
Biotinylated Human DKK1 C terminal Domain Protein

Cat. No. DKK-HM51CB

ᠺ᠕ᡗ᠋ᠴᡃ᠐ᠫ Description Recombinant Biotinylated Human DKK1 C terminal Domain Protein is expressed from HEK293 with hFc tag and Avi tag at the C-Terminus. Source It contains Met178-His266. Accession O94907 Molecular The protein has a predicted MW of 38.69 kDa. Due to glycosylation, the protein migrates to 50-55 kDa based on Weight Bis-Tris PAGE result. Endotoxin Less than 1 EU per µg by the LAL method. Purity > 95% as determined by Bis-Tris PAGE Formulation and Storage Lyophilized from 0.22µm filtered solution in 20mM NaAc, 150mM NaCl (pH 5.0). Normally 8% trehalose is added Formulation as protectant before lyophilization. Dissolve the lyophilized protein in 20mM NaAc, 150mM NaCl (pH 5.0). Please refer to the Certificate of Analysis Reconstitution for detailed instructions. -20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend Storage to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. Background Dickkopf-1 (Dkk1), the founding and best-studied member of the Dkk family, functions as an antagonist of canonical Wnt/β-catenin. Dkk1 is considered to play a broad role in a variety of biological processes.

Assay Data

Bis-Tris PAGE




Biotinylated Human DKK1 C terminal Domain on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

Biotinylated Human DKK1 C terminal Domain Protein

Cat. No. DKK-HM51CB

Assay Data

Biotinylated DKK1 C terminal Domain, hFc Tag ELISA 0.5µg Human LRP-6, mFc Tag Per Well



Immobilized Human LRP-6, mFc Tag at 5μ g/ml (100 μ l/well) on the plate. Dose response curve for Biotinylated DKK1 C terminal Domain, hFc Tag with the EC50 of 2.07 μ g/ml determined by ELISA (QC Test).


KVCJUS

Log Biotinylated DKK1 C terminal Domain, hFc Tag Conc.($\mu g/ml)$

Biotinylated Human DKK1 C terminal Domain, hFc Tag ELISA

ELISA Data

For Research Use Only



Immobilized Anti-DKK1 Antibody, hFc Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Human DKK1 C terminal Domain, hFc Tag with the EC50 of 19.7ng/ml determined by ELISA (QC Test).