

Human DKK1 C terminal Domain Protein



Cat. No.     DKK-HM51C

Description	
Source	Recombinant Human DKK1 C terminal Domain Protein is expressed from HEK293 with hFc tag and Avi tag at the C-Terminus. It contains Met178-His266.
Accession	O94907
Molecular Weight	The protein has a predicted MW of 38.69 kDa. Due to glycosylation, the protein migrates to 45-55 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE


Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in 20mM NaAc,150mM NaCl (pH 5.0). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in 20mM NaAc,150mM NaCl (pH 5.0). Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

**Background**

Dickkopf-1 (Dkk1), the founding and best-studied member of the Dkk family, functions as an antagonist of canonical Wnt/β-catenin. Dkk1 is considered to play a broad role in a variety of biological processes.

Assay Data

Bis-Tris PAGE




Human DKK1 C terminal Domain on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

ELISA Data

Assay Data

Human DKK1 C terminal Domain, hFc Tag ELISA

0.05µg Human DKK1 C terminal Domain, hFc Tag Per Well



Immobilized Human DKK1 C terminal Domain, hFc Tag at 0.5 µg/ml (100 µl/Well) on the plate. Dose response curve for Biotinylated Anti-DKK1 Antibody, hFc Tag with the EC50 of 26.7 ng/ml determined by ELISA.