# Biotinylated Human DR3/TNFRSF25 Protein

### Cat. No. DR3-HM401B



Description	
Source	Recombinant Biotinylated Human DR3/TNFRSF25 Protein is expressed from HEK293 with His tag and Avi tag at the C-terminus.
	It contains Gln25-Gln199.
Accession	AAI17190.1
Molecular Weight	The protein has a predicted MW of 26.40 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

## Formulation and Storage

Formulation Supplied as 0.22 µm filtered solution in PBS (pH 7.4).

Storage Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller


quantities for optimal storage. Please minimize freeze-thaw cycles.

## **Background**

Death Receptor 3 (DR3), also known as TNFRSF25, TRAMP, LARD, or WSL-1, is a death-domain-containing TNF-family receptor that, like its closest paralog TNFR1, binds the adaptor molecule TRADD through its cytoplasmic death domain. TRADD recruitment endows DR3 with dual-signaling capability to activate NF-κB and MAP-kinase signaling or alternatively trigger caspase activation and programmed cell death.

## **Assay Data**

### **Bis-Tris PAGE**



**ELISA Data** 


Biotinylated Human DR3 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

## **Assay Data**



# Biotinylated Human DR3, His-Avi Tag ELISA

0.1μg Biotinylated Human DR3, His-Avi Tag Per Well



Log Human TNFSF15, hFc Tag Conc.(μg/ml)

Immobilized Biotinylated Human DR3, His-Avi Tag at  $1\mu g/ml$  ( $100\mu l/well$ ) on the streptavidin precoated plate ( $5\mu g/ml$ ). Dose response curve for Human TNFSF15, hFc Tag with the EC50 of 15.2ng/ml determined by ELISA (QC Test).