

Canine EGF Protein

Cat. No. EGF-DE101




Description	
Source	Recombinant Canine EGF Protein is expressed from E.coli with His tag at the N-terminus. It contains Asn973-Arg1024.
Accession	Q9BEA0
Molecular Weight	The protein has a predicted MW of 13.61 kDa. The protein migrates to 16-22 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
The epidermal growth factor (EGF) family of peptides encodes several proteins that can function as growth factors. The EGF-like peptides, with the exception of proteins of the EGF-CFC subfamily, bind and activate tyrosine kinase receptors that belong to the erbB family.	

Assay Data


Bis-Tris PAGE



Canine EGF on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC


Assay Data



The purity of Canine EGF is greater than 95% as determined by SEC-HPLC.

ELISA Data


Canine EGF, His Tag ELISA
0.1µg Canine EGF, His Tag Per Well



Immobilized Canine EGF, His Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Human EGFR, hFc Tag with the EC50 of 7.0ng/ml determined by ELISA (QC Test).

Cell Based Assay

Recombinant Canine EGF Bioactivity



Measured in a cell proliferation assay using Balb 3T3 mouse embryonic fibroblasts. The EC50 for this effect is typically 0.5 - 3 ng/mL.