

Biotinylated Human EGFR/HER1 Protein



Cat. No. EGF-HM401B

Description	
Source	Recombinant Biotinylated Human EGFR/HER1 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Leu25-Ser645.
Accession	P00533-1
Molecular Weight	The protein has a predicted MW of 71.5 kDa. Due to glycosylation, the protein migrates to 90-120 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
The epidermal growth factor receptor is a transmembrane protein that is a receptor for members of the epidermal growth factor family of extracellular protein ligands. The epidermal growth factor receptor is a member of the ErbB family of receptors, a subfamily of four closely related receptor tyrosine kinases: EGFR, HER2/neu, Her 3 and Her 4. Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses.	

Assay Data

Bis-Tris PAGE



Biotinylated Human EGFR on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Biotinylated Human EGFR/HER1 Protein

Cat. No. EGF-HM401B




Assay Data



The purity of Biotinylated Human EGFR is greater than 95% as determined by SEC-HPLC.


ELISA Data

Biotinylated Human EGFR, His Tag ELISA
0.1µg Biotinylated Human EGFR, His Tag Per Well



Immobilized Biotinylated Human EGFR, His Tag at 1µg/ml (100µl/well) on the streptavidin precoated plate(5µg/ml). Dose response curve for Human EGF, hFc Tag with the EC50 of 4.1ng/ml determined by ELISA (QC Test).

SPR Data



Biotinylated Human EGFR, His Tag captured on CM5 Chip via Anti-his antibody can bind Human EGF, No Tag with an affinity constant of 0.62 nM as determined in SPR assay (Biacore T200).