Canine EMMPRIN/CD147 Protein

Cat. No. EPN-DM147



Recombinant Canine EMMPRIN/CD147 Protein is expressed from HEK293 with His tag at the C-Terminus.
It contains Ala22-Arg205.
XP_533964.2
The protein has a predicted MW of 21.54 kDa. Due to glycosylation, the protein migrates to 30-45 kDa based on Bis-Tris PAGE result.
Less than 1 EU per μg by the LAL method.
> 95% as determined by Bis-Tris PAGE
> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

CD147, also known as extracellular matrix metalloproteinase inducer (EMMPRIN) or basigin, is expressed in a variety of cell types. It is involved in the regulation of extracellular matrix (ECM) remodeling during physiological and pathological processes including wound healing, inflammatory diseases, and cancer. CD147 is a diagnostic and therapeutic target in cancer and inflammatory diseases, either directly or indirectly, by targeting CD147 partners.

Assay Data

Bis-Tris PAGE



Canine EMMPRIN on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Canine EMMPRIN/CD147 Protein

Cat. No. EPN-DM147



Assay Data



The purity of Canine EMMPRIN is greater than 95% as determined by SEC-HPLC.