

Human EMMPRIN/CD147 Protein



Cat. No. EPN-HM147


Description	
Source	Recombinant Human EMMPRIN/CD147 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Ala22-His205.
Accession	NP_940991.1
Molecular Weight	The protein has a predicted MW of 21.2 kDa. Due to glycosylation, the protein migrates to 28-40 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
CD147, also known as extracellular matrix metalloproteinase inducer (EMMPRIN) or basigin, is expressed in a variety of cell types. It is involved in the regulation of extracellular matrix (ECM) remodeling during physiological and pathological processes including wound healing, inflammatory diseases, and cancer. CD147 is a diagnostic and therapeutic target in cancer and inflammatory diseases, either directly or indirectly, by targeting CD147 partners.	

Assay Data


Bis-Tris PAGE



Human EMMPRIN on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Human EMMPRIN is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human EMMPRIN, His Tag ELISA

0.1µg Human EMMPRIN, His Tag Per Well



Immobilized Human EMMPRIN, His Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Anti-EMMPRIN Antibody, hFc Tag with the EC50 of 3.0ng/ml determined by ELISA (QC Test).