

## Mouse EREG Protein

Cat. No. ERE-MM201



### Description

Source	Recombinant Mouse EREG Protein is expressed from HEK293 with hFc tag at the N-Terminus.
	It contains Val56-Leu101.
Accession	Q61521
Molecular Weight	The protein has a predicted MW of 32.7 kDa. Due to glycosylation, the protein migrates to 37-40 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

### Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

### Background

Epidermal growth factor receptor (EGFR) regulates many crucial cellular programs, with seven different activating ligands shaping cell signaling in distinct ways. EGFR ligands epiregulin (EREG) and epigen (EPGN) stabilize different dimeric conformations of the EGFR extracellular region. As a consequence, EREG or EPGN induce less stable EGFR dimers than EGF-making them partial agonists of EGFR dimerization. Unexpectedly, this weakened dimerization elicits more sustained EGFR signaling than seen with EGF, provoking responses in breast cancer cells associated with differentiation rather than proliferation.

### Assay Data

#### Bis-Tris PAGE



Mouse EREG on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


#### SEC-HPLC

## Mouse EREG Protein

Cat. No. ERE-MM201



### Assay Data



The purity of Mouse EREG is greater than 95% as determined by SEC-HPLC.