

Mouse FLT3 Ligand Protein



Cat. No. FLT-MM13L


Description	
Source	Recombinant Mouse FLT3 Ligand Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Gly27-Gln189.
Accession	P49772-1
Molecular Weight	The protein has a predicted MW of 19.61 kDa. Due to glycosylation, the protein migrates to 28-38 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
Flt3 Ligand, also known as FL, is an alpha -helical cytokine that promotes the differentiation of multiple hematopoietic cell lineages. Stimulates the proliferation of early hematopoietic cells by activating FLT3. Synergizes well with a number of other colony stimulating factors and interleukins.	

Assay Data

Bis-Tris PAGE



Mouse FLT3 Ligand on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Mouse FLT3 Ligand Protein

Cat. No. FLT-MM13L




Assay Data



The purity of Mouse FLT3 Ligand is greater than 95% as determined by SEC-HPLC.


ELISA Data

Mouse FLT3 Ligand, His Tag ELISA
0.2μg Mouse FLT3 Ligand, His Tag Per Well



Immobilized Mouse FLT3 Ligand, His Tag at 2μg/ml (100μl/well) on the plate. Dose response curve for Human FLT3, hFc Tag with the EC50 of 20.0ng/ml determined by ELISA (QC Test).

SPR Data



Mouse FLT3 Ligand, His Tag captured on CM5 Chip via anti-his antibody can bind Human FLT3, hFc Tag with an affinity constant of 0.18 nM as determined in SPR assay (Biacore T200).