Human GCGR/Glucagon receptor Protein

Cat. No. GCR-HM40R



Description	
Source	Recombinant Human GCGR/Glucagon receptor Protein is expressed from HEK293 with His tag and Avi tag at the C-terminus.
	It contains Ala26-Lys136.
Accession	P47871
Molecular Weight	The protein has a predicted MW of 15.96 kDa. Due to glycosylation, the protein migrates to 30-50 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	The pancreatic hormone glucagon activates the glucagon receptor (GCGR), a class B seven-transmembrane G protein-coupled receptor that couples to the stimulatory heterotrimeric G protein and provokes PKA-dependent


signaling cascades vital to hepatic glucose metabolism and islet insulin secretion. The GCGR expressed at the

plasma membrane is constitutively ubiquitinated and upon agonist-activation, internalized GCGRs are

deubiquitinated at early endosomes and recycled via Rab4-containing vesicles.

Assay Data

Bis-Tris PAGE




SEC-HPLC

Human GCGR on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

Cat. No. GCR-HM40R

KAGTUS

Assay Data



The purity of Human GCGR is greater than 95% as determined by SEC-HPLC.