Mouse HGFA Protein (pro form)

Cat. No. HGF-MM10A



| Description | |
|---------------------|--|
| Source | Recombinant Mouse HGFA Protein (pro form) is expressed from HEK293 with His tag at the C-Terminus. |
| | It contains Gln35-Ser653. |
| Accession | Q9R098 |
| Molecular Weight | The protein has a predicted MW of 67.9 kDa. Due to glycosylation, the protein (pro form) migrates to 80-100 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1 EU per μg by the LAL method. |
| Purity | > 95% as determined by Bis-Tris PAGE |
| | > 95% as determined by HPLC |
| | |

Formulation and Storage


| Formulation | Lyophilized from 0.22µm filtered solution in 20mM Tris,150mM NaCl, 2mM CaCl2 (pH 8.0). Normally 8% trehalose is added as protectant before lyophilization. |
|----------------|---|
| Reconstitution | Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

Hepatocyte growth factor activator (HGFA) is a serine protease initially identified as a potent activator of hepatocyte growth factor/scatter factor. Hepatocyte growth factor/scatter factor is known to be critically involved in tissue morphogenesis, regeneration, and tumor progression, via its receptor, MET.In vivo, HGFA also activates macrophage-stimulating protein, which has roles in macrophage recruitment and inflammatory processes, cellular survival and wound healing through its receptor, RON.

Assay Data

Bis-Tris PAGE




Mouse HGFA (pro form) on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cat. No. HGF-MM10A



Assay Data



The purity of Mouse HGFA (pro form) is greater than 95% as determined by SEC-HPLC.