

Human ICAM-3/CD50 Protein



Cat. No. IAM-HM103


Description	
Source	Recombinant Human ICAM-3/CD50 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Gln30-His485.
Accession	CAA49369.1
Molecular Weight	The protein has a predicted MW of 50.54 kDa. Due to glycosylation, the protein migrates to 80-110 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
ICAM-3 interacts with LFA1, and is involved in the intercellular adhesion of leukocytes as well as in the mainenance of cell survival. ICAM-3 expression induces cancer cell proliferation, and an increase in ICAM-3 expression can contribute to cancer progression.	

Assay Data

Bis-Tris PAGE



Human ICAM-3 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.