# Human IL-15RA/IL-15 R alpha/CD215 Protein


Cat. No. ILR-HM215



Cat. 140. ILIX-I IIVIZ 13	
Description	
Source	Recombinant Human IL-15RA/IL-15 R alpha/CD215 Protein is expressed from HEK293 with hFc tag at the C-Terminus.
	It contains Ile31-Thr205.
Accession	Q13261-1
Molecular Weight	The protein has a predicted MW of 45.1 kDa. Due to glycosylation, the protein migrates to 70-80 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	Interleukin-15 receptor alpha (IL-15R alpha) is a high affinity IL-15 binding protein that is crucial for mediating IL-15 functions such as memory CD8 T cell proliferation and NK, NK/T cell, and intestinal intraepithelial lymphocyte development.

## **Assay Data**

### **Bis-Tris PAGE**




Human IL-15RA on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

**SEC-HPLC** 

# KAGTUS


### **Assay Data**



The purity of Human IL-15RA is greater than 95% as determined by SEC-HPLC.

#### **ELISA Data**

Human IL-15RA, hFc Tag ELISA 0.2μg Human IL-15, No Tag Per Well



Immobilized Human IL-15, No Tag at 2µg/ml (100µl/Well) on the plate. Dose response curve for Human IL-15RA, hFc Tag with the EC50 of 24.0ng/ml determined by ELISA (QC Test).