Biotinylated Human IL-5 R alpha/CD125 Protein

Cat. No. ILR-HM45RB




Description	
Source	Recombinant Biotinylated Human IL-5 R alpha/CD125 Protein is expressed from HEK293 with His tag and Avitag at the C-Terminus.
	It contains Asp21-Glu335.
Accession	Q01344-1
Molecular Weight	The protein has a predicted MW of 38.7 kDa. Due to glycosylation, the protein migrates to 55-60 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per ug by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 24 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	

Interleukin5 Receptor alpha (IL5 R alpha), also known as CD125, is a 60 kDa hematopoietin receptor that plays a

dominant role in eosinophil biology. This is the receptor for interleukin-5. The alpha chain binds to IL5.

Assay Data

Bis-Tris PAGE




Biotinylated Human IL-5 R alpha on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

KAGTUS

Assay Data




The purity of Biotinylated Human IL-5 R alpha is greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated Human IL-5 R alpha, His Tag ELISA

0.2μg Human IL-5, His Tag Per Well



Log Biotinylated Human IL-5 R alpha, His Tag Conc.(μg/ml)

Immobilized Human IL-5, His Tag at $2\mu g/ml$ (100 μ I/well) on the plate. Dose response curve for Biotinylated Human IL-5 R alpha, His Tag with the EC50 of $0.15\mu g/ml$ determined by ELISA.