

Human Integrin alpha M beta 2 (ITGAM&ITGB2) Heterodimer Protein



Cat. No. ITG-HM1MB


Description	
Source	Recombinant Human Integrin alpha M beta 2 (ITGAM&ITGB2) Heterodimer Protein is expressed from HEK293 with His tag at the C-Terminus of ITGAM It contains Phe17-Asn1104 (ITGAM) acidic tail and Gln23-Asn700 (ITGB2) basic tail.
Accession	P11215-1(ITGAM)&P05107-1(ITGB2)
Molecular Weight	The protein has a predicted MW of 126.10 kDa (ITGAM) and 79.50 kDa (ITGB2). Due to glycosylation, the protein migrates to 140-160 kDa (ITGAM) and 90-100 kDa (ITGB2) based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
Integrin αMβ2, also called Mac-1 or complement receptor-3 (CR3), is a heterodimer expressed on monocytes. It functions as both an adhesion molecule mediating the diapedesis of leukocytes across the endothelium and a receptor for the iC3b fragment of complement responsible for phagocytic/degranulation responses to microorganisms[1]. Active αMβ2 binds iC3b, coagulation proteins fibrinogen, plasminogen and factor X, extracellular matrix (ECM) proteins fibronectin, laminin and collagen, and cell surface ICAMs, myelin basic protein and DCSIGN[2].	

Assay Data

Bis-Tris PAGE




Human ITGAM&ITGB2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

ELISA Data

Human ITGAM&ITGB2, His Tag ELISA

0.2µg Human ITGAM&ITGB2, His Tag Per Well



Immobilized Human ITGAM&ITGB2, His Tag at 2 µg/ml (100 µl/well) on the plate. Dose response curve for Anti-ITGB2 Antibody, hFc Tag with the EC50 of 11.9 ng/ml determined by ELISA (QC Test).