

Biotinylated Mouse Integrin alpha V beta 3 (ITGAV&ITGB3) Heterodimer Protein



Cat. No. ITG-MM4V3B


Description	
Source	Recombinant Biotinylated Mouse Integrin alpha V beta 3 (ITGAV&ITGB3) Heterodimer Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus of ITGAV. It contains Phe31-Val988(ITGAV) acidic tail and Glu26-Asp717(ITGB3) basic tail.
Accession	P43406(ITGAV)&O54890(ITGB3)
Molecular Weight	The protein has a predicted MW of 108.77 kDa (ITGAV) & 80.84 kDa (ITGB3). Due to glycosylation, the protein migrates to 90-130 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
ITGAV&ITGB3 binds to NRG1 (via EGF domain) and this binding is essential for NRG1-ERBB signaling. ITGAV&ITGB3 binds to FGF1 and this binding is essential for FGF1 signaling. ITGAV&ITGB3 binds to FGF2 and this binding is essential for FGF2 signaling. ITGAV&ITGB3 binds to IGF1 and this binding is essential for IGF1 signaling. ITGAV&ITGB3 binds to IGF2 and this binding is essential for IGF2 signaling.	

Assay Data

Bis-Tris PAGE




Biotinylated Mouse ITGAV&ITGB3 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

ELISA Data

Biotinylated Mouse ITGAV&ITGB3, His Tag ELISA

0.5µg Human Vitronectin, His Tag Per Well



Immobilized Human Vitronectin, His Tag at 5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Mouse ITGAV&ITGB3, His Tag with the EC50 of 3.2µg/ml determined by ELISA.