

Human LILRB2/CD85d/ILT4 Protein

Cat. No. LIL-HM3B2



Description

Source	Recombinant Human LILRB2/CD85d/ILT4 Protein is expressed from HEK293 with mFc (IgG1) tag at the C-Terminus.
	It contains Gln22-Arg457.
Accession	AAL36991.1
Molecular Weight	The protein has a predicted MW of 73.6 kDa. Due to glycosylation, the protein migrates to 80-110 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The immunoglobulin-like transcript (ILT) comprise a family of activating and inhibitory type immunoreceptors whose genes are located in the same locus that encodes killer cell Ig-like receptors (KIR). ILT4, also known as LIR-2 and LILRB2, is a type I transmembrane protein expressed primarily on monocytes and dendritic cells (DC). LILRB2 is a receptor for class I MHC antigens. Recognizes a broad spectrum of HLA-A, HLA-B, HLA-C, HLA-G and HLA-F alleles.

Assay Data

Bis-Tris PAGE




Human LILRB2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

ELISA Data

Human LILRB2, mFc Tag ELISA

0.05µg Human LILRB2, mFc Tag Per Well




Immobilized Human LILRB2, mFc Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Anti-LILRB2 Antibody, hFc Tag with the EC50 of 13.0ng/ml determined by ELISA (QC Test).

Assay Data

Blocking Data

Inhibition of Human LILRB2 and HLA-G Binding
0.2µg Human LILRB2, mFc Tag Per Well



Serial dilutions of Anti-LILRB2 Antibody were added into Human HLA-G Complex Tetramer, His Tag : Human LILRB2, mFc Tag binding reactions. The half maximal inhibitory concentration (IC50) is 0.11µg/ml.