

Cynomolgus MCP-1/CCL2 Protein

Cat. No. MCP-CM101

Description

Source	Recombinant Cynomolgus MCP-1/CCL2 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Gln24-Pro99.
Accession	P61274
Molecular Weight	The protein has a predicted MW of 9.79 kDa. Due to glycosylation, the protein migrates to 13-16 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μ g by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 μ m filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Monocyte chemoattractant protein-1 (MCP-1) (also referred to as chemokine (C-C motif) ligand 2 (CCL2)) is expressed by mainly inflammatory cells and endothelial cells. MCP-1 has been reported to play an important role in the pathogenesis of atherosclerosis and considerable evidence supports that the monocyte containing MCPs and macrophage influences the growth of other cell types within the atherosclerotic lesion. This review will focus on the general structure features of MCP-1 and its role in atherosclerosis.

Assay Data


Bis-Tris PAGE



Cynomolgus MCP-1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Cynomolgus MCP-1 is greater than 95% as determined by SEC-HPLC.