#### Chimeric HLA-A\*02:01 (mα3) &B2M&WT-1 (RMFPNAPYL) Tetramer Protein





| Description         |  |
|---------------------|--|
| Source              | Recombinant Chimeric HLA-A*02:01(mα3)&B2M&WT-1 (RMFPNAPYL) Tetramer Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus, tetramer is assembled by biotinylated monomer and streptavidin. |
|                     | It contains Gly25-Thr206(Human HLA-A*02:01 $\alpha$ 1& $\alpha$ 2)&Asp207-Glu299(Mouse H-2Ld $\alpha$ 3), Ile21-Met119 (B2M) and RMFPNAPYL peptide.  |
| Accession           | A0A140T913(Human HLA-A*02:01 α1&α2)&P01897(Mouse H-2Ld α3)&P61769(B2M)&RMFPNAPYL   |
| Molecular<br>Weight | The protein has a predicted MW of 258 kDa. Due to glycosylation, the protein migrates to 260-265 kDa under Non reducing (N) condition based on Bis-Tris PAGE result.   |
| Endotoxin           | Less than 1 EU per ug by the LAL method.   |
| Purity              | > 95% as determined by Bis-Tris PAGE   |
|                     | > 95% as determined by HPLC  |
| Formulation and     | l Storage  |
| Formulation         | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.   |

Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed

# Reconstitution

-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend

to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.


# **Background**

Storage

The WT1 protein plays a role in cell growth, the process by which cells mature to perform specific functions (differentiation), and the self-destruction of cells (apoptosis). WT1 is differentially expressed in serous, endometrioid, clear cell, and mucinous carcinomas of the peritoneum, fallopian tube, ovary, and endometrium. The Human HLA-A\*0201 WT-1 (RMFPNAPYL) complex Protein is a complex of HLA-A\*0201 of the MHC Class I, B2M and RMFPNAPYL peptide of the WT-1.

### **Assay Data**

## **Bis-Tris PAGE**




Chimeric HLA-A\*02:01 (ma3) &B2M&WT-1 (RMFPNAPYL) Tetramer on Bis-Tris PAGE under Non reducing (N) condition. The purity is greater than 95%.

**SEC-HPLC** 



#### **Assay Data**




The purity of Chimeric HLA-A\*02:01 (m $\alpha$ 3) &B2M&WT-1 (RMFPNAPYL) Tetramer is greater than 95% as determined by SEC-HPLC.

#### **ELISA Data**

# Chimeric HLA-A\*02:01(m $\alpha$ 3)&B2M&WT-1 Tetramer, His Tag ELISA

0.2μg Chimeric HLA-A\*02:01(mα3)&B2M&WT-1 Tetramer, His Tag Per Well



Log Anti-HLA-A\*02:01&B2M&WT-1 Antibody, hFc Tag Conc.(µg/ml)

Immobilized Chimeric HLA-A\*02:01 (m $\alpha$ 3) &B2M&WT-1 (RMFPNAPYL) Tetramer, His Tag at 2 $\mu$ g/ml (100 $\mu$ l/well) on the plate. Dose response curve for Anti-HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Antibody, hFc Tag with the EC50 of 6.3ng/ml determined by ELISA.