# Biotinylated Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Monomer Protein






Out. 110. 111110 11111110	
Description	
Source	Recombinant Biotinylated Human HLA-A*02:01&B2M&WT-1 (RMFPNAPYL) Monomer Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Gly25-Thr305(HLA-A*02:01), Ile21-Met119(B2M) and RMFPNAPYL peptide.
Accession	A0A140T913(HLA-A*02:01)&P61769(B2M)&RMFPNAPYL
Molecular Weight	The protein has a predicted MW of 50.5 kDa. Due to glycosylation, the protein migrates to 55-60 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	The WT1 protein plays a role in cell growth, the process by which cells mature to perform specific functions (differentiation), and the self-destruction of cells (apoptosis). WT1 is differentially expressed in serous, endometrioid, clear cell, and mucinous carcinomas of the peritoneum, fallopian tube, ovary, and endometrium. The Human HLA-A*0201 WT-1 (RMFPNAPYL) complex Protein is a complex of HLA-A*0201 of

the MHC Class I, B2M and RMFPNAPYL peptide of the WT-1.

# **Assay Data**

### **Bis-Tris PAGE**




Biotinylated Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Monomer on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

**SEC-HPLC** 




### **Assay Data**



The purity of Biotinylated Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Monomer was greater than 95% as determined by SEC-HPLC.

### **ELISA Data**

Biotinylated Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Monomer, His-Avi Tag ELISA 0.1µg Biotinylated Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Monomer, His-Avi Tag Per Well



Immobilized Biotinylated Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Monomer, His Tag at 1µg/ml (100µl/Well) on streptavidin (5µg/ml) precoated plate. Dose response curve for Anti-HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Antibody, hFc Tag with the EC50 of 6.7ng/ml determined by ELISA.