## Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Monomer Protein






Oat. 140. WII 10-1 IIVI-13	'
Description	
Source	Recombinant Human HLA-A*02:01&B2M&WT-1 (RMFPNAPYL) Monomer Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Gly25-Thr305(HLA-A*02:01), Ile21-Met119(B2M) and RMFPNAPYL peptide.
Accession	A0A140T913(HLA-A*02:01)&P61769(B2M)&RMFPNAPYL
Molecular Weight	The protein has a predicted MW of 50.5 kDa. Due to glycosylation, the protein migrates to 52-60 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	The WT1 protein plays a role in cell growth, the process by which cells mature to perform specific functions (differentiation), and the self-destruction of cells (apoptosis). WT1 is differentially expressed in serous, endometrioid, clear cell, and mucinous carcinomas of the peritoneum, fallopian tube, ovary, and endometrium. The Human HLA-A*0201 WT-1 (RMFPNAPYL) complex Protein is a complex of HLA-A*0201 of

the MHC Class I, B2M and RMFPNAPYL peptide of the WT-1.

# **Assay Data**

### **Bis-Tris PAGE**



Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Monomer on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


**SEC-HPLC** 

### Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Monomer Protein

Cat. No. MHC-HM431



## **Assay Data**




The purity of Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Monomer was greater than 95% as determined by SEC-HPLC.

#### **ELISA Data**

### Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL), His Tag ELISA


0.2μg Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL), His Tag Per Well



Log Anti-HLA-A\*02:01&B2M&WT-1 Antibody, hFc Tag Conc.(µg/ml)

Immobilized Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Monomer, His Tag at 2µg/ml (100µl/well) on the plate. Dose response curve for Anti-HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Antibody, hFc Tag with the EC50 of 19.0ng/ml determined by ELISA (QC Test).

#### **SPR Data**



Human HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Monomer, His Tag captured on CM5 Chip via Anti-His Antibody can bind Anti-HLA-A\*02:01&B2M&WT-1 (RMFPNAPYL) Antibody, hFc Tag with an affinity constant of 0.33nM as determined in SPR assay (Biacore T200).