## Mouse CD10/MME Protein

Cat. No. MME-MM110



Description	
Source	Recombinant Mouse CD10/MME Protein is expressed from HEK293 with His tag at the N-Terminus.
	It contains Tyr52-Trp750.
Accession	Q61391
Molecular Weight	The protein has a predicted MW of 81.1 kDa. Due to glycosylation, the protein migrates to 82-110 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

## Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

# **Background**

CD10 is an endopeptidase that degrades various bioactive peptides in the extracellular matrix. In addition to enzymatic degradation, it affects multiple intracellular signal transduction pathways. CD10 expression has been extensively studied in human epithelial cancers of numerous organs and sites. CD10 expression pattern depended on the histotypes of thyroid lesions. When possible metastatic tumours and non-epithelial tumours are excluded, high CD10 expression may be useful in determining whether a primary thyroid carcinoma includes an anaplastic component.

## **Assay Data**

## **Bis-Tris PAGE**



Mouse CD10 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


**SEC-HPLC** 

# Mouse CD10/MME Protein

Cat. No. MME-MM110

# KAGTUS

# **Assay Data**



The purity of Mouse CD10 is greater than 95% as determined by SEC-HPLC.  $\label{eq:cd2} % \begin{tabular}{ll} \end{tabular} % \begin{tab$ 

## **Bioactivity Data**

Measured by its ability to cleave the fluorogenic peptide substrate, Mca-RPPGFSAFK (Dnp) -OH. The specific activity is > 3000 pmoles/min/µg.