Human MSLN/Mesothelin Protein

Cat. No. MSL-HM28D



Description	
Source	Recombinant Human MSLN/Mesothelin Protein is expressed from HEK293 with hFc tag at the C-Terminus.
	It contains Glu296-Arg606.
Accession	Q13421-2
Molecular Weight	The protein has a predicted MW of 61.54 kDa. Due to glycosylation, the protein migrates to 68-78 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage

Formulation Supplied as 0.22µm filtered solution in PBS (pH 7.4).

Storage Valid for 12 months from date of receipt when stored at -80°C.Recommend to aliquot the protein into smaller


quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Mesothelin, also known as MSLN, is a protein that in humans is encoded by the MSLN gene. Cloning studies showed that the mesothelin gene encodes a precursor protein that is processed to yield mesothelin which is attached to the cell membrane by a glycophosphatidylinositol linkage and a 31-kDa shed fragment named megakaryocyte-potentiating factor (MPF). Although it has been proposed that mesothelin may be involved in cell adhesion, its biological function is not known. A knockout mouse line that lacks mesothelin reproduces and develops normally.

Assay Data


Bis-Tris PAGE



Human MSLN on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

KAGTUS


Assay Data



The purity of Human MSLN is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human MSLN, hFc Tag ELISA 0.05μg Human MSLN, hFc Tag Per Well



Log Biotinylated Anti-MSLN Antibody, hFc Tag Conc.(µg/ml)

Immobilized Human MSLN, hFc Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Biotinylated Anti-MSLN Antibody, hFc Tag with the EC50 of 21.5ng/ml determined by ELISA.