

Human NPC2 Protein



Cat. No. NPC-HM102

Description	
Source	Recombinant Human NPC2 Protein is expressed from HEK293 with His tag at the C-terminus. It contains Glu20-Leu151.
Accession	P61916-1
Molecular Weight	The protein has a predicted MW of 15.67 kDa. Due to glycosylation, the protein migrates to 21-24 kDa and 27-35 kDa kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
The Niemann Pick type C (NPC) proteins, NPC1 and NPC2, are involved in the lysosomal storage disease, NPC disease. The formation of a NPC1NPC2 proteinprotein complex is believed to be necessary for the transfer of cholesterol and lipids out of the late endosomal (LE)/lysosomal (Lys) compartments. Mutations in either NPC1 or NPC2 can lead to an accumulation of cholesterol and lipids in the LE/Lys, the primary phenotype of the NPC disease.	

Assay Data

Bis-Tris PAGE



Human NPC2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.