

Mouse Oncostatin M/OSM Protein



Cat. No. OSM-MM102


Description	
Source	Recombinant Mouse Oncostatin M/OSM Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Asn25-Arg206.
Accession	P53347
Molecular Weight	The protein has a predicted MW of 21.64 kDa. Due to glycosylation, the protein migrates to 35-45 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
OSM is a pleiotropic cytokine that initiates its biological activities by binding to specific cell surface receptors. Inhibits the proliferation of a number of tumor cell lines. Stimulates proliferation of AIDS-KS cells. It regulates cytokine production, including IL-6, G-CSF and GM-CSF from endothelial cells. Uses both type I OSM receptor (heterodimers composed of LIFR and IL6ST) and type II OSM receptor (heterodimers composed of OSMR and IL6ST). Involved in the maturation of fetal hepatocytes, thereby promoting liver development and regeneration.	

Assay Data

Bis-Tris PAGE



Mouse Oncostatin M on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Mouse Oncostatin M/OSM Protein

Cat. No. OSM-MM102



Assay Data



The purity of Mouse Oncostatin M is greater than 95% as determined by SEC-HPLC.