### Biotinylated Human PSMA/FOLH1 Protein

### Cat. No. PSM-HM410B




Description	
Source	Recombinant Biotinylated Human PSMA/FOLH1 Protein is expressed from HEK293 with His tag and Avi tag at the N-terminus.
	It contains Lys44-Ala750.
Accession	Q04609-1
Molecular Weight	The protein has a predicted MW of 82.43 kDa. Due to glycosylation, the protein migrates to 95-115 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 90% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22 $\mu$ m filtered solution in 25 mM MES, 500 mM NaCl (pH 6.5). Normally 8% mannitol is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	

Prostate-specific membrane antigen (PSMA) is an enzyme that in humans is encoded by the FOLH1 (folate hydrolase 1) gene, also known as Glutamate carboxypeptidase II (GCPII). Human PSMA is highly expressed in the prostate, roughly a hundred times greater than in most other tissues. In some prostate cancers, PSMA is the second-most upregulated gene product, with an 8- to 12-fold increase over levels in noncancerous prostate cells.

### **Assay Data**

### **Bis-Tris PAGE**




Biotinylated Human PSMA on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

**SEC-HPLC** 

# KAGTUS

### **Assay Data**




The purity of Biotinylated Human PSMA is greater than 90% as determined by SEC-HPLC.

### **ELISA Data**

## Biotinylated Human PSMA, His-Avi Tag ELISA

0.1μg Biotinylated Human PSMA, His-Avi Tag Per Well



 $Log\ Anti-PSMA\ Antibody,\ hFc\ Tag\ Conc.(\mu g/ml)$ 

Immobilized Biotinylated Human PSMA, His-Avi Tag at 1µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Anti-PSMA Antibody, hFc Tag with the EC50 of 4.1ng/ml determined by ELISA (QC Test).