

Human RETN Protein

Cat. No. RET-HM201



Description

Source	Recombinant Human RETN Protein is expressed from HEK293 with hFc tag at the N-Terminus.
	It contains Ser17-Pro108.
Accession	Q9HD89-1
Molecular Weight	The protein has a predicted MW of 36.9 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Resistin (RETN) is a hormone secreted by adipocytes, which plays an important role in glucose and lipid metabolism. The RETN gene is expressed in goat various analyzed tissues, and the results showed that the expression of RETN gene in lung tissue was higher than that in other analyzed tissues of goat ($p < .01$). Moreover, the expression level of RETN gene in the goat's intramuscular preadipocytes decreased first and then increased, and reached the highest on the fifth day, which was significantly higher than that of undifferentiated intramuscular preadipocytes.

Assay Data


Bis-Tris PAGE



Human RETN on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Human RETN is greater than 95% as determined by SEC-HPLC.