

Human Siglec-3/CD33 Protein

Cat. No. SIG-HM203



Description

Source	Recombinant Human Siglec-3/CD33 Protein is expressed from HEK293 with hFc tag at the C-Terminus.
	It contains Asp18-His259.
Accession	P20138-1
Molecular Weight	The protein has a predicted MW of 53.6 kDa. Due to glycosylation, the protein migrates to 68-80 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Sialic-acid-binding immunoglobulin-like lectin (Siglec) that plays a role in mediating cell-cell interactions and in maintaining immune cells in a resting state. They are sialoadhesin/CD169/Siglec-1, CD22/Siglec-2, CD33/Siglec-3, Myelin-Associated Glycoprotein (MAG/Siglec-4a) and Siglecs 5 to 11. To date, no Siglec has been shown to recognized any cell surface ligand other than sialic acids, suggesting that interactions with glycans containing this carbohydrate are important in mediating the biological functions of Siglecs.

Assay Data


Bis-Tris PAGE



Human Siglec-3 on Bis-Tris PAGE under reduced conditions. The purity is greater than 95%.

SEC-HPLC


Assay Data



The purity of Human Siglec-3 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human Siglec-3, hFc Tag ELISA
0.05µg Human Siglec-3, hFc Tag Per Well



Immobilized Human Siglec-3, hFc Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Anti-Siglec-3 Antibody, hFc Tag with the EC50 of 21.8ng/ml determined by ELISA.