Human CD48/SLAMF2 Protein

Cat. No. SLA-HM2MF



Description	
Source	Recombinant Human CD48/SLAMF2 Protein is expressed from HEK293 with hFc tag at the C-Terminus.
	It contains Gln27-Ser220.
Accession	P09326-1
Molecular Weight	The protein has a predicted MW of 48.9 kDa. Due to glycosylation, the protein migrates to 62-72 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and	Storage


Formulation	Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

CD48, also known as BLAST-1, BCM-1, and SLAMF2, is a 65 kDa GPI-linked protein in the CD2 family of immunoglobulin superfamily molecules.CD2 and 2B4 (CD244) are known ligands for CD48. CD48 protein is expressed on most lineage-committed hematopoietic cells but not on hematopoietic stem cells or multipotent hematopoietic progenitors.

Assay Data

Bis-Tris PAGE




Human CD48 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

KAGTUS


Assay Data



The purity of Human CD48 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human CD48, hFc Tag ELISA 0.1µg Human 2B4, His Tag Per Well



Immobilized Human 2B4, His Tag at $1\mu g/ml$ (100 $\mu l/well$) on the plate. Dose response curve for Human CD48, hFc Tag with the EC50 of 0.34 $\mu g/ml$ determined by ELISA.