

Biotinylated Human SIRP gamma/CD172g Protein



Cat. No. SRP-HM40GB


Description	
Source	Recombinant Biotinylated Human SIRP gamma/CD172g Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Glu29-Pro360.
Accession	Q9P1W8
Molecular Weight	The protein has a predicted MW of 39.7 kDa. Due to glycosylation, the protein migrates to 48-53 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
Signal regulatory protein gamma (SIRP gamma, designated CD172g), also called SIRP beta 2, is a monomeric 45-47 kDa type I transmembrane protein belonging to the SIRP/SHPS (CD172) family of the Ig superfamily.SIRP gamma is probable immunoglobulin-like cell surface receptor. On binding with CD47, mediates cell-cell adhesion. Engagement on T-cells by CD47 on antigen-presenting cells results in enhanced antigen-specific T-cell proliferation and costimulates T-cell activation.	

Assay Data

Bis-Tris PAGE



Biotinylated Human SIRP gamma on Bis-Tris PAGE under reduced condition. The purity is greater than 95%..


SEC-HPLC

Biotinylated Human SIRP gamma/CD172g Protein

Cat. No. SRP-HM40GB



Assay Data




The purity of Biotinylated Human SIRP gamma is greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated Human SIRP gamma, His Tag ELISA

0.2µg Biotinylated Human SIRP gamma, His Tag Per Well



Immobilized Biotinylated Human SIRP Gamma, His Tag at 2µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Human CD47, hFc Tag with the EC50 of 0.28µg/ml determined by ELISA.