

Biotinylated Human SIRP alpha V2/CD172a Protein



Cat. No.    SRP-HM4V2B


Description	
Source	Recombinant Biotinylated Human SIRP alpha V2/CD172a Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Glu31-Arg369.
Accession	AAH26692.1
Molecular Weight	The protein has a predicted MW of 40 kDa. Due to glycosylation, the protein migrates to 55-65 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per ug by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
Signal regulatory protein α (SIRPα) is a regulatory membrane glycoprotein from SIRP family expressed mainly by myeloid cells and also by stem cells or neurons.SIRPα acts as inhibitory receptor and interacts with a broadly expressed transmembrane protein CD47 also called the "don't eat me" signal.Cancer cells highly expressed CD47 that activate SIRP α and inhibit macrophage-mediated destruction.	

Assay Data

Bis-Tris PAGE



Biotinylated Human SIRP alpha V2/CD172a on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Biotinylated Human SIRP alpha V2/CD172a Protein

Cat. No.    SRP-HM4V2B



Assay Data




The purity of Biotinylated Human SIRP alpha V2/CD172a is greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated Human SIRP alpha V2, His Tag ELISA

0.2µg Biotinylated Human SIRP alpha V2, His Tag Per Well



Immobilized Biotinylated Human SIRP alpha V2, His Tag at 2µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Human CD47, hFc Tag with the EC50 of 2.5ng/ml determined by ELISA (QC Test).