

Biotinylated Human VEGF-C/Flt4-L Protein



Cat. No. VEG-HM4F1B


Description	
Source	Recombinant Biotinylated Human VEGF-C/Flt4-L Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Thr103-Arg227.
Accession	Q6FH59
Molecular Weight	The protein has a predicted MW of 17.1 kDa. Due to glycosylation, the protein migrates to 23-30 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in 50mM MES, 150mM NaCl (pH 6.0). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in 50mM MES, 150mM NaCl (pH 6.0). Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
the lymphangiogenic factors vascular endothelial growth factor C (VEGFC) and VEGFD are cleaved by thrombin and plasmin, serine proteases generated during hemostasis and wound healing. Genetic studies reveal that platelet enhancement of lymphatic growth after wounding is dependent on the release of VEGFC, but not VEGFD, a finding consistent with high expression of VEGFC in both platelets and avian thrombocytes.	

Assay Data

Bis-Tris PAGE




Biotinylated Human VEGF-C on Bis-Tris PAGE under reduced (R) condition. The purity is greater than 95%.

ELISA Data

Biotinylated Human VEGF-C, His Tag ELISA

0.1µg Biotinylated Human VEGF-C, His Tag Per Well



Immobilized Biotinylated Human VEGF-C, His Tag at 1µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Human VEGF R3, hFc Tag with the EC50 of 8.2ng/ml determined by ELISA.