

Biotinylated Human VEGF R2/KDR Protein (Primary Amine Labeling)



Cat. No. VGF-HM3R2B


Description	
Source	Recombinant Biotinylated Human VEGF R2/KDR Protein (Primary Amine Labeling) is expressed from HEK293 with mFc (IgG1) tag at the C-Terminus. It contains Ala20-Glu764.
Accession	P35968-1
Molecular Weight	The protein has a predicted MW of 110 kDa. Due to glycosylation, the protein migrates to 135-160 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
Tyrosine-protein kinase that acts as a cell-surface receptor for VEGFA, VEGFB and PGF, and plays an essential role in the development of embryonic vasculature, the regulation of angiogenesis, cell survival, cell migration, macrophage function, chemotaxis, and cancer cell invasion. The tyrosine kinase receptor vascular endothelial growth factor receptor 2 (VEGFR2) is a key regulator of angiogenesis.	

Assay Data

Bis-Tris PAGE



Biotinylated Human VEGF R2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Biotinylated Human VEGF R2/KDR Protein (Primary Amine Labeling)



Cat. No. VGF-HM3R2B

Assay Data




The purity of Biotinylated Human VEGF R2 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated Human VEGF R2, mFc Tag ELISA

0.5µg Human VEGF165, No Tag Per Well



Immobilized Human VEGF165, No Tag at 5µg/ml (100µl/Well) on the plate. Dose response curve for Biotinylated Human VEGF R2, mFc Tag with the EC50 of 98.8ng/ml determined by ELISA.