# NKMAXBio We support you, we believe in your research

# **Human beta-II Tubulin antibody**

Catalog Number: ATGA0196

### **PRODUCT INFORMATION**

#### Catalog number

ATGA0196

#### Clone No.

AT5B2

#### **Product type**

Monoclonal Antibody

#### UnitProt No.

Q9BVA1

#### **NCBI Accession No.**

NP 821080

#### **Alternative Names**

TUBB2B, Tubulin beta 2B class lib, Tubulin beta 2b

### **PRODUCT SPECIFICATION**

#### **Antibody Host**

Mouse

#### **Reacts With**

Human

#### Concentration

1mg/ml (determined by BCA assay)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

#### **Immunogen**

Recombinant human beta-Tubulin (1-445aa) purified from E. coli

## Isotype

IgG2b kappa

#### **Purification Note**

By protein-G affinity chromatography

### **Application**

ELISA, WB, ICC/IF

#### Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

#### **Storage**



# NKMAXBio We support you, we believe in your research

## **Human beta-II Tubulin antibody**

Catalog Number: ATGA0196

Can be stored at +4C For long term storage, aliquot and store at -20C. Avoid repeated freezing and thawing cycles.

#### **BACKGROUND**

### **Description**


Tubulin is a major cytoskeleton component that has five distinct forms. Multiple beta Tubulin isoforms have been characterized and are expressed in mammalian tissues. Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain.

#### **General References**

Xavier Hubert Jaglin, et al. (2009) Nat Genet, 41(6):746-52. Bouwmeester T, et al. (2004) Nat Cell Bio, 6(2):97-105. Modig, C, et al. (1999) Cell Motil Cytoskeleton, 42(4): 315-330.

#### **DATA**

#### Western blot analysis (WB)




The cell lysates(40ug) and tissue lysates(35ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human beta-Tubulin (1:5000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: Jurkat cell lysate Lane 2.: A549 cell lysate Lane 3.: A431 cell lysate Lane 4.: HepG2 cell lysate

Lane 5.: Mouse brain tissue lyate

#### Immunocytochemistry/Immunofluorescence (ICC/IF)




Immunofluorescence of HaCaT cells stained with Hoechst 3342 (Blue) for nucleus staining and monoclonal anti-human beta-Tubulin antibody (1:1000) with Texas Red (Red).




# NKMAXBio We support you, we believe in your research

# **Human beta-II Tubulin antibody**

Catalog Number: ATGA0196



Immunofluorescence of A431 cells stained with DAPI (Blue) for nucleus staining and monoclonal anti-human beta-Tubulin antibody (1:500) with Alexa Fluor 488 (Green).



Immunofluorescence of Balb/3T3 cells stained with DAPI (Blue) for nucleus staining and monoclonal anti-human beta-Tubulin antibody (1:200) with Alexa Fluor 488 (Green).

