

Human SOD1/Cu-Zn SOD antibody

Catalog Number: ATGA0405

PRODUCT INFORMATION

Catalog number

ATGA0405

Clone No.

AT6F1

Product type

Monoclonal Antibody

UnitProt No.

P00441

NCBI Accession No.

NP_000445

Alternative Names

Superoxide dismutase 1 soluble, ALS, SOD, ALS1, IPOA, Cu-Zn superoxide dismutase, Superoxide dismutase 1 soluble, SOD1, Superoxide dismutase 1, soluble ALS 1, Amyotrophic lateral sclerosis 1 Amyotrophic lateral sclerosis 1 adult, Cu/Zn SOD, Cu/Zn superoxide dismutase, Homodimer, Indophenoloxidase A, Mn superoxide dismutase, SOD 1, SOD soluble, SOD2, SODC, Superoxide dismutase 1 soluble, Superoxide dismutase Cu Zn, Superoxide dismutase cystolic

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human SOD1 (1-154aa) purified from E. coli

Isotype

IgG1 kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, ICC/IF, FACS

Usage

The antibody has been tested by ELISA, Western blot, ICC/IF and FACS analysis to assure specificity and

Human SOD1/Cu-Zn SOD antibody

Catalog Number: ATGA0405

reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


Cu/Zn superoxide dismutase (SOD1), is a major antioxidant enzyme that catalyzes the conversion of superoxide anion to hydrogen peroxide and molecular oxygen. SOD1 facilitates the dismutation of oxygen radicals to hydrogen peroxide and also catalyzes pro-oxidant reactions, which include the peroxidase activity and hydroxyl radical generating activity. Also over 100 distinct SOD1 inherited mutations have been identified in the familial form of amyotrophic lateral sclerosis (ALS), a progressive degenerative disease of motor neurons. The mechanism by which mutant SOD1 induces the neurodegeneration observed in ALS is still unclear.

General References

Deng HX., et al. (1993) Science. 261(5124): 1047-51.
Valentine JS., et al. (2003) Proc Natl Acad Sci USA. 100(7): 3617-22.
Rosen DR., et al. (1993) Nature. 362(6415): 59-62.


DATA

Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human SOD1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1. : A549 cell lysate
Lane 2. : A431 cell lysate
Lane 3. : 293T cell lysate
Lane 4. : PC3 cell lysate
Lane 5. : U87MG cell lysate




The recombinant proteins (20ng) of SOD two isoform were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human SOD1 (1:1000).

Lane 1 : Recombinant protein SOD1
Lane 2 : Recombinant protein SOD2

Human SOD1/Cu-Zn SOD antibody


Catalog Number: ATGA0405

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of SOD1 in Hep3B cells. The cell was stained with ATGA0405 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

Flow cytometry (FACS)



Flow cytometry analysis of SOD1 in Hep3B cell line, staining at 2-5ug for 1×10^6 cells (red line). The secondary antibody used goat anti-mouse IgG Alexa fluor 488 conjugate. Isotype control antibody was mouse IgG (black line).