NKMAXBio We support you, we believe in your research

Human Serpin E1/PAI-1 antibody

Catalog Number: ATGA0458

PRODUCT INFORMATION

Catalog number

ATGA0458

Clone No.

AT1C5

Product type

Monoclonal Antibody

UnitProt No.

P05121

NCBI Accession No.

NP 000593

Alternative Names

Plasminogen activator inhibitor-1, PLANH1, SERPINE1, PAI1, PLANH1, Plasminogen activator inhibitor-1, Plasminogen activator inhibitor-1

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human PAI1 (24-402aa) purified from E. coli.

Isotype

IgG1 kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot analysis and ICC/IF to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBio We support you, we believe in your research

Human Serpin E1/PAI-1 antibody

Catalog Number: ATGA0458

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


Plasminogen activator inhibitor-1 (PAI-1), a member of the serine protease inhibitor (serpin) super family. PAI-1 is the principal inhibitor of tissue plasminogen activator and urokinase, the activators of plasminogen and hence fibrinolysis. PAI-1 is mainly produced by the endothelium but is also secreted by other tissue types, such as adipose tissue.

General References

Gorlatova NV., et al.(2007) J Biol Chem. 282(12): 9288-9296. Renckens R., et al. (2006) J Immunol. 177(11): 8171-8176.

DATA

Western blot analysis (WB)




The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human PAI1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: 293T cell lysate

Lane 2.: PAI1 transfected 293T cell lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of PAI1 in HeLa cells. The cell was stained with ATGA0458 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

