# **PRODUCT INFORMATION**

Expression system E.coli

**Domain** 1-235aa

**UniProt No.** Q9NQT4

NCBI Accession No. NP\_064543.3

**Alternative Names** Exosome complex component RRP46, hRrp46p, p12B, RRP41B, RRP46, Rrp46p

# **PRODUCT SPECIFICATION**

**Molecular Weight** 27.5 kDa (256aa) confirmed by MALDI-TOF

### **Concentration** 0.5mg/ml (determined by Bradford assay)

### Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 2mM DTT, 30% glycerol, 200mM NaCl

### Purity

> 90% by SDS-PAGE

**Tag** His-Tag

Application SDS-PAGE

### **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## BACKGROUND

### Description

EXOSC5, also known as exosome component 5, belongs to the RNase PH family. EXOSC5 has been shown to interact with Exosome component 8 and Exosome component 1. It is Non-catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease activity and participates in a multitude of cellular RNA processing and degradation events. It highly expressed in a variety of hematopoietic and epithelial tumor cell lines, but not in normal hematopoietic tissues or other normal tissue, with the exception of testis. Recombinant human EXOSC5 protein, fused to His-tag at N-terminus, was expressed in E. coli and purified by using



conventional chromatography techniques.

#### **Amino acid Sequence**

<MGSSHHHHHH SSGLVPRGSH M>MEEETHTDA KIRAENGTGS SPRGPGCSLR HFACEQNLLS RPDGSASFLQ GDTSVLAGVY GPAEVKVSKE IFNKATLEVI LRPKIGLPGV AEKSRERLIR NTCEAVVLGT LHPRTSITVV LQVVSDAGSL LACCLNAACM ALVDAGVPMR ALFCGVACAL DSDGTLVLDP TSKQEKEARA VLTFALDSVE RKLLMSSTKG LYSDTELQQC LAAAQAASQH VFRFYRESLQ RRYSKS

### **General References**

Mukherjee D., et al. (2002) EMBO J. 21:165-174 Basu u., et al. (2011) Cell 144:353-363

## DATA

#### **SDS-PAGE**



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.