PRODUCT INFORMATION

Expression system E.coli

Domain 29-146aa

UniProt No. P12273

NCBI Accession No. NP_002643

Alternative Names Prolactin-inducible protein, GCDFP-15, GCDFP15, GPIP4

PRODUCT SPECIFICATION

Molecular Weight 15.9 kDa (141aa)

Concentration 1mg/ml (determined by Bradford assay)

Formulation

Liquid in. 20mM Tris-HCl buffer (pH 8.0) containing 0.4M urea, 10% glycerol

Purity

> 90% by SDS-PAGE

Tag His-Tag

Application SDS-PAGE, Denatured

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

Prolactin-inducible protein, also known as PIP, is a major protein component of benign breast gross cysts. It is a known marker of breast cancer, as it is found in approximately 50% of all breast cancer specimens. It is expressed in exocrine glands and, in pathologic conditions, in breast cysts and breast cancers exhibiting apocrine features. PIP and prostate specific antigen are co-expressed in androgen receptor-positive breast tumours. Recombinant human PIP protein, fused to His-tag at N-terminus, was expressed in E. coli.



Amino acid Sequence

MGSSHHHHHH SSGLVPRGSH MGSQDNTRKI IIKNFDIPKS VRPNDEVTAV LAVQTELKEC MVVKTYLISS IPLQGAFNYK YTACLCDDNP KTFYWDFYTN RTVQIAAVVD VIRELGICPD DAAVIPIKNN RFYTIEILKV E

General References

Mazoujian G., et al. (1983) J Pathol. 110:105-112.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

15% SDS-PAGE (3ug)