PRODUCT INFORMATION

Expression system E.coli

Domain 76-590aa

UniProt No. Q9BQG1

NCBI Accession No. NP_115674

Alternative Names Synaptotagmin-3, Synaptotagmin-3, SytIII

PRODUCT SPECIFICATION

Molecular Weight 57.7 kDa (540aa)

Concentration 1mg/ml (determined by Bradford assay)

Formulation

Liquid in. 20mM Tris-HCl buffer (pH 8.0) containing 10% glycerol 0.4M urea

Purity

> 85% by SDS-PAGE

Tag His-Tag

Application SDS-PAGE,Denatured

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

SYT3, also known as synaptotagmin-3, belongs to the synaptotagmin family. It may be involved in Ca (2+) dependent exocytosis of secretory vesicles through Ca (2+) and phospholipid binding to the C2 domain or may serve as Ca (2+) sensors in the process of vesicular trafficking and exocytosis. Recombinant human SYT3 protein, fused to His-tag at N-terminus, was expressed in E. coli.

Amino acid Sequence

MGSSHHHHHH SSGLVPRGSH MGSEFWKLCW VPWRDKGGSA VGGGPLRKDL GPGVGLAGLV GGGGHHLAAG



LGGHPLLGGP HHHAHAAHHP PFAELLEPGS LGGSDTPEPS YLDMDSYPEA AAAAVAAGVK PSQTSPELPS EGGAGSGLLL LPPSGGGLPS AQSHQQVTSL APTTRYPALP RPLTQQTLTS QPDPSSEERP PALPLPLPGG EEKAKLIGQI KPELYQGTGP GGRRSGGGPG SGEAGTGAPC GRISFALRYL YGSDQLVVRI LQALDLPAKD SNGFSDPYVK IYLLPDRKKK FQTKVHRKTL NPVFNETFQF SVPLAELAQR KLHFSVYDFD RFSRHDLIGQ VVLDNLLELA EQPPDRPLWR DIVEGGSEKA DLGELNFSLC YLPTAGRLTV TIIKASNLKA MDLTGFSDPY VKASLISEGR RLKKRKTSIK KNTLNPTYNE ALVFDVAPES VENVGLSIAV VDYDCIGHNE VIGVCRVGPD AADPHGREHW AEMLANPRKP VEHWHQLVEE KTVTSFTKGS KGLSEKENSE

General References

Jones JM. et al. (1995) Mamm Genome. 6:212-213. Li C. et al. (1995) Nature. 375:594-599.

DATA



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

15% SDS-PAGE (3ug)

