# **PRODUCT INFORMATION**

**Expression system** E.coli

**Domain** 45-374aa

**UniProt No.** Q9BY67

NCBI Accession No. NP\_055148

### **Alternative Names**

Cell adhesion molecule 1, Immunoglobulin superfamily member 4, IgSF4, Nectin-like protein 2, NECL-2, Spermatogenic immunoglobulin superfamily, SgIgSF, Synaptic cell adhesion molecule, SynCAM, Tumor suppressor in lung cancer 1, TSLC-1, ST17, BL2, IGSF4A, SYNCAM1, RA175

# **PRODUCT SPECIFICATION**

### **Molecular Weight**

39.4 kDa (353aa) confirmed by MALDI-TOF

#### Concentration

0.25mg/ml (determined by Bradford assay)

#### Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol, 1mM DTT

Purity

> 85% by SDS-PAGE

Tag

His-Tag

Application SDS-PAGE

#### **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## BACKGROUND

## Description

CADM1 mediates homophilic cell-cell adhesion in a Ca2+-independent manner and also mediates heterophilic cell-cell adhesion with CADM3 and PVRL3 in a Ca2+-independent manner. This protein acts as a tumor suppressor in non-small-cell lung cancer (NSCLC) cells. Interaction with CRTAM promotes natural killer (NK) cell cytotoxicity and interferon-gamma (IFN-gamma) secretion by CD8+ cells in vitro as well as NK cell-mediated



rejection of tumors expressing CADM3 in vivo. Recombinant human CADM1 protein, fused to His-tag at N-terminus, was expressed in E. coli and purified by using conventional chromatography techniques.

#### Amino acid Sequence

#### **General References**

Barton C.H., et al. (1988) Development 104:165-173.

## DATA



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

15% SDS-PAGE (3ug)

