# **PRODUCT INFORMATION**

**Expression system** E.coli

**Domain** 311-426aa

**UniProt No.** P08476

NCBI Accession No. NP\_002183

Alternative Names Inhibin beta A chain, EDF, FRP

# **PRODUCT SPECIFICATION**

Molecular Weight 15.2 kDa (137aa)

**Concentration** 1mg/ml (determined by Bradford assay)

### Formulation

Liquid in. 20mM Tris-HCl buffer (pH 8.0) containing 10% glycerol

**Purity** > 90% by SDS-PAGE

**Tag** His-Tag

Application SDS-PAGE, Denatured

## **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## BACKGROUND

### Description

INHBA also known as inhibin beta A chain. The inhibin beta A subunit joins the alpha subunit to form a pituitary FSH secretion inhibitor. Inhibin has been shown to regulate gonadal stromal cell proliferation negatively and to have tumor-suppressor activity. In addition, serum levels of inhibin have been shown to reflect the size of granulosa-cell tumors and can therefore be used as a marker for primary as well as recurrent disease. Recombinant human INHBA, fused to His-tag at N-terminus, was expressed in E. coli.



### Amino acid Sequence

<MGSSHHHHHH SSGLVPRGSH M>GLECDGKVN ICCKKQFFVS FKDIGWNDWI IAPSGYHANY CEGECPSHIA GTSGSSLSFH STVINHYRMR GHSPFANLKS CCVPTKLRPM SMLYYDDGQN IIKKDIQNMI VEECGCS

### **General References**

Antsiferova M., et al. (2013) J. Immunol. 191 (12), 6147-6155 Woeckel VJ., et al. (2013) J. Cell. Physiol. 228 (11), 2167-2174

