PRODUCT INFORMATION

Expression system Baculovirus

Domain 21-140aa

UniProt No. P21460

NCBI Accession No. NP_034106

Alternative Names CST3, CysC

PRODUCT SPECIFICATION

Molecular Weight 14.2 kDa (126aa) confirmed by MALDI-TOF

Concentration 1mg/ml (determined by absorbance at 280nm)

Formulation Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity > 95% by SDS-PAGE

Endotoxin level < 1 EU per 1ug of protein (determined by LAL method)

Tag His-Tag

Application SDS-PAGE

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

CST3, also known as Cystatin-C is a member of family 2 of the cystatin superfamily. This protein is a secreted type 2 cysteine protease inhibitor synthesized in all nucleated cells, has been proposed as a replacement for serum creatinine for the assessment of renal function, particularly to detect small reductions in glomerular filtration rate. It is involved in processes such as tumor invasion and metastasis, inflammation and some



neurological diseases. It inhibits many cysteine proteases such as papain and cathepsins B, H, K, L and S. It indicates that a murine model should be relevant for studies of the human disease, hereditary Cystatin C amyloid angiopathy. Recombinant Mouse CST3, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Amino acid Sequence

ATPKQGPRML GAPEEADANE EGVRRALDFA VSEYNKGSND AYHSRAIQVV RARKQLVAGV NYFLDVEMGR TTCTKSQTNL TDCPFHDQPH LMRKALCSFQ IYSVPWKGTH SLTKFSCKNA <HHHHHH>

General References

Janowski. R., et al. (2001) Nat. Struct. Biol. 8(4):316-20. Mussap M., et al. (2004) Crit Rev Clin Lab Sci. 41(5-6):467-550.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

