# NKMAXBio We support you, we believe in your research

## Recombinant human C1q R1/CD93 protein

Catalog Number: ATGP3395

## **PRODUCT INFORMATION**

## **Expression system**

Baculovirus

#### **Domain**

22-580aa

#### UniProt No.

O9NPY3

#### **NCBI Accession No.**

NP 036204

#### **Alternative Names**

Complement component C1q receptor, CD93, C1qR(P), C1QR1, C1qRP, CDw93, dJ737E23.1, ECSM3, MXRA4, C1q receptor 1, C1q/MBL/SPA receptor, C1qR, C1qR(p), C1qr1, C1QR1\_HUMAN, C1qRp, CD93, CD93 antigen, CD93 molecule, CDw93, Complement component 1 q subcomponent receptor 1, Complement component C1q receptor, dJ737E23.1, ECSM3, Matrix remodeling associated protein 4, Matrix remodelling associated 4, Matrix-remodeling-associated protein 4, MXRA4

## **PRODUCT SPECIFICATION**

## **Molecular Weight**

59.3 kDa (567aa)

#### Concentration

0.5mg/ml (determined by absorbance at 280nm)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

#### **Purity**

> 90% by SDS-PAGE

## **Endotoxin level**

< 1 EU per 1ug of protein (determined by LAL method)

#### Tag

His-Tag

## **Application**

SDS-PAGE

## **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## **BACKGROUND**



## Recombinant human C1q R1/CD93 protein

Catalog Number: ATGP3395

## **Description**

CD93, also known as complement component C1q receptor, belongs to a family of transmembrane glycoproteins which also includes endosialin and thrombomodulin. It is involved in various aspects of inflammatory reactions. CD93 is expressed by vascular endothelial cells and by a variety of hematopoietic cells. It is receptor for C1q, mannose-binding lectin (MBL2) and pulmonary surfactant protein A (SPA). Recombinant human CD93, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.


#### **Amino acid Sequence**

TGADTEAVVC VGTACYTAHS GKLSAAEAQN HCNQNGGNLA TVKSKEEAQH VQRVLAQLLR REAALTARMS KFWIGLQREK GKCLDPSLPL KGFSWVGGGE DTPYSNWHKE LRNSCISKRC VSLLLDLSQP LLPSRLPKWS EGPCGSPGSP GSNIEGFVCK FSFKGMCRPL ALGGPGQVTY TTPFQTTSSS LEAVPFASAA NVACGEGDKD ETQSHYFLCK EKAPDVFDWG SSGPLCVSPK YGCNFNNGGC HQDCFEGGDG SFLCGCRPGF RLLDDLVTCA SRNPCSSSPC RGGATCVLGP HGKNYTCRCP QGYQLDSSQL DCVDVDECQD SPCAQECVNT PGGFRCECWV GYEPGGPGEG ACQDVDECAL GRSPCAQGCT NTDGSFHCSC EEGYVLAGED GTQCQDVDEC VGPGGPLCDS LCFNTQGSFH CGCLPGWVLA PNGVSCTMGP VSLGPPSGPP DEEDKGEKEG STVPRAATAS PTRGPEGTPK ATPTTSRPSL SSDAPITSAP LKMLAPSGSP GVWREPSIHH ATAASGPQEP AGGDSSVATQ NNDGTDGQKV EHHHHHHH

#### **General References**

McGreal EP., et al. (2002) J Immunol. 168:5222-5232. Chevrier S., et al. (2009) Proc Natl Acad Sci USA. 106:3895-3900.

## **DATA**



15% SDS-PAGE (3ug)-

3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

